

Noah Zinkin, MD, MMSc
775 Park Avenue, Suite 225
Huntington, NY 11743
(631) 923-1420

Moviprep – Evening Before Prep

*Please remember to arrange for a responsible adult to accompany you home. If you do not have an adult escort your procedure will be cancelled and rescheduled. Taxis are not permitted.

Review the preparation schedule below for the days preceding your colonoscopy. Should you need further assistance please call 631-923-1420

1 Week Prior

Go to pharmacy and fill your prescription for MoviPrep. Please call our office if you did not receive a prescription.

If you take fiber supplements or vitamins containing iron, please discontinue 5 days before your appointment.

If you have diabetes, please ask us for diet and medication instructions.

If you take aspirin, plavix, Coumadin, or other blood thinners, please discuss with Dr. Zinkin whether you should continue or discontinue these prior to your procedure.

3 Days Prior

Begin Low Fiber Diet (No fruits or vegetables. No whole wheat or high fiber. No seeds, nuts, corn or popcorn No bran or bulking agents)

The Day Prior

You may have a light, low fiber breakfast. Afterwards, begin a clear liquid diet. This includes water, clear broth or bouillon, coffee or tea, Gatorade, carbonated & non-carbonated soft drinks, Kool-Aid or other fruit-flavored drinks, strained fruit juices (no pulp), Jell-O, popsicles and hard candy. For lunch, you may have ice cream, a milkshake, or yogurt that does not contain seeds, nuts or anything solid. Drink at least 64 ounces (1/2 gallon) of clear liquid throughout the day.

NO RED LIQUIDS

MoviPrep Instructions:

*You may mix the solution ahead of time and refrigerate prior to drinking. The solution once mixed must be used within 24 hours.

Empty 1 pouch A and 1 pouch B into the disposable container

Add a clear liquid (we recommend lemonade or unsweetened iced tea) to the top line of the container. Mix to dissolve.

Starting at 5:30 PM the evening before your procedure:

The MoviPrep container is divided by 4 marks. Every 15 minutes drink the solution down to the next mark (approximately 8 ounces), until the full liter has been completed. Drink additional half liter (16 ounces) of clear liquid over the next hour.

Starting at 10 PM the evening before your procedure:

Drink the second liter of solution following the same process completed the earlier: Every 15 minutes drink the solution down to the next mark (approximately 8 ounces) until the liter has been completed. Drink additional half liter (16 ounces) of clear liquid over the next hour.

Do not eat or drink anything after midnight

On the procedure day:

Do not eat or drink anything. Morning medications may be taken with a sip of water.

Leave your valuables at home

Please arrange for a responsible driver to take you home. Taxis are not permitted.

Remember that no driving or work is permitted on the day of the procedure.

Women who have had a period within 1 year will be asked for a urine specimen for a pregnancy test