

Noah Zinkin, MD, MMSc
775 Park Avenue, Suite 225
Huntington, NY 11743
(631) 923-1420

Suprep – Split Dose Prep

*Please remember to arrange for a responsible adult to accompany you home. If you do not have an adult escort your procedure will be cancelled and rescheduled. Taxis are not permitted.

Review the preparation schedule below for the days preceding your colonoscopy. Should you need further assistance please call 631-923-1420

1 Week Prior

Go to pharmacy and fill your prescription for Suprep. Please call our office if you did not receive a prescription.

If you take fiber supplements or vitamins containing iron, please discontinue 5 days before your appointment.

If you have diabetes, please ask us for diet and medication instructions.

If you take aspirin, plavix, Coumadin, or other blood thinners, please discuss with Dr. Zinkin whether you should continue or discontinue these prior to your procedure.

3 Days Prior

Begin Low Fiber Diet (No raw fruits or vegetables, no whole wheat or high fiber, no seeds, nuts or popcorn, no bran or bulking agents)

The Day Prior

You may have a light, low fiber breakfast. Afterwards, begin a clear liquid diet. This includes water, clear broth or bouillon, coffee or tea, Gatorade, carbonated & non-carbonated soft drinks, Kool-Aid or other fruit-flavored drinks, strained fruit juices (no pulp), Jell-O, popsicles and hard candy. For lunch, you may have ice cream, a milkshake, or yogurt that does not contain seeds, nuts or anything solid. Drink at least 64 ounces (1/2 gallon) of clear liquid throughout the day.

NO RED LIQUIDS

SUPREP Instructions:

Starting at 5:30 PM the evening before your procedure:

1. Pour ONE six ounce bottle of SUPREP into the mixing container
2. Add cool drinking water to the 16 ounce line on the container and mix
3. Drink all the liquid in the container
4. You must drink 2 or more 16 ounce containers of water over the next 1 hour

On the procedure day

Starting 6 hours before your procedure:

1. Pour the second six ounce bottle of SUPREP into the mixing container
2. Add cool drinking water to the 16 ounce line on the container and mix
3. Drink all the liquid in the container
4. You must drink 2 or more 16 ounce containers of water over the next 1 hour

Have NOTHING to eat or drink for 4 hours before your procedure. Morning medications may be taken with a sip of water.

Upon departing for your procedure:

Do not eat or drink anything. Morning medications may be taken with a sip of water.

Leave your valuables at home

Please arrange for a responsible driver to take you home. Taxis are not permitted.

Remember that no driving or work is permitted on the day of the procedure.